

Journal of Sociology & Cultural Research Review (JSCRR)

Available Online: <https://jscrr.edu.com.pk>

Print ISSN: [3007-3103](#) Online ISSN: [3007-3111](#)

Platform & Workflow by: [Open Journal Systems](#)

PREVALENCE OF HAMSTRING TIGHTNESS DUE TO EMOTIONAL SIGNIFICANCE IN BODYBUILDERS OF ISLAMABAD

Farwa Shafique

Clinical & Educational Psychologist, Department of Psychology, International Islamic University, Islamabad

Hassan Imran

PhD Scholar, Department of Psychology, Riphah International University Faisalabad Campus

hassanimran332@gmail.com

Sidrah Shaheen

Assistant Professor Psychology, HED, KP

ABSTRACT

Hamstring tightness is not unusual problem in humans because of their running patterns or posturing and shortage of sports. The bodybuilders who have low probabilities of emotional distress they have lower hamstring tightness that lower the dangers for musculoskeletal muscle injuries. Muscle tightness can also cause by emotional traumatic experiences happened in past. Human body collect these traumas in mind and create tension. Stressful events in life cause emotional disturbance which cause tension in mind and body also. The purpose of the study is to determine the prevalence of hamstring tightness in bodybuilders of Islamabad who got emotional stress. This study became executed as cross-sectional examine in Islamabad. The variety of pattern become 377 bodybuilders in Islamabad. From the entire wide variety of 377 bodybuilder 82% of the candidate had been determined as having occurrence of hamstring tightness from which 3.01% had been mendacity in underweight, 82% have been every day, 10% have been approximately 25 BMI and 4.5% have been above 30 BMI with extra obese. The least hamstring tightness changed into tested in bodybuilders in Islamabad. Standardized equipment have been used with .99 reliability, musculoskeletal fitness questionnaire with .90 reliability and emotional scale with .92 reliability. Only the better weight humans had better tendency of hamstring tightness who had good BMI and lower stress by releasing old tensions.

Keywords: emotions, hamstring tightness, musculoskeletal fitness, lower extremity

Introduction

Prevalence of hamstring tightness is found in nearly all around the global mainly in athletes (Weerasekara, et al. 2013). Ergonomics is the study of different posture and positions during work (Akinpelu et al, 2009). Ergonomics came into existence in 1949 Oxford, England when there was a group of people whom work was related to human performance came together and held a meeting in which it was decided to develop a new word for this field called ergonomics. It is a Greek word 'ergo' means for work and 'nomos' means natural laws. The concept of ergonomics was given by Paul Fitts in which he describes that human performance are based on five factors Learning, individual differences, motivation, environment and task.

Hamstring tightness can be prevented by controlling the emotions. When person experiences a particular painful emotional experience. It show effects on the body

of individuals by releasing chemical products in brain. Due to each complex interconnection body and brain controls the muscle movement combine (Koli & Anap, 2018). Some people deals with the poisoned emotions based on their sensitivity. This toxic emotion physically demonstrate imbalances in our body. Emotions rather than consciously moving the body. High incidence of hamstring tightness some of the college students of university of 18-25 age is present (Fatima et al, 2017). Hamstring tightness occurs in many of the college students is due to extended sitting (Bhavana et al, 2013). If the individual have a strong anxiety, the imbalance can lead to allergic reactions. An unconstrained toxic mood will produce chronic pain.

“Inability to increase the knee absolutely while the hip is flexed observed through pain or ache alongside the posterior thigh and/or knee is commonly attributed to hamstring muscle tightness” (Mistry, Vyas and Sheth, 2013). Hamstring tightness happens in early life and it has a tendency to boom with age. The revolutionary decline in flexibility with age has been attributed to alternate in elasticity and reduced stage of bodily sports (Gleim & McHugh, 1997). Tight hamstring may make any pose difficult.

The weight lifting and lumbar pelvic rhythm reason hamstring tightness. Tightness reason the shortening of muscle which lower the variety of lively or passive mobility. Muscle maximum prone to tightness in decrease extremity are hamstring muscle mass (Mayorga-Vega et al, 2015). Emotional reasons and their influence shall make a lot of sense. It is almost like a map of all chemicals flow.

Hamstring extensibility is a physical fitness component widely recognized as an important marker of health and quality of life (Erkula et al, 2002). Hamstring extensibility is a bodily health issue broadly diagnosed as a critical marker of fitness and pleasant of life (Raether & Lutter, 1982). Safe, relaxation posture is ideal for stress relief.

Stress is a major problem among shoe manufacturing workers. The prevalence of labor-related depression was relatively high among shoe cobblers and also there is less support from the society with no work protection. Workers suffer discomfort and pain as there is increase in exertion and the healing or rehabilitation time is less (Shah &, Iftakhar, 2017). Thing can be improve reducing work related demand and providing protection to the workers (Esola et al, 1974).

Hamstring tightness in male and woman is different. Male have greater hamstring tightness in left leg whilst lady have in right leg due to loss of bodily sports (Dionne et al, 1974)). Fitness to keep extreme pressure on the body and intense use of muscle can cause damage of muscles. Mostly athletes work on their fitness level during development phase. They continue to strengthen their muscles which can cause muscle tightness (Sahrmann, 2002)

Previous researches showed that poor BMI can cause the higher hamstring muscle tightness directly. They showed lower BMI and high obesity. It also can be occur due to the less exercises and physical activities, who have lazy behavior. So the researcher became encouraged for doing study about that how is the prevalence of hamstring muscle tightness with lower extremity functions and musculoskeletal health. Previous researches showed that the body builders got high hamstring tightness especially males more than females. They used to do exercises on daily basis. If body builders use precautions and maintain BMI they will adjustable in

health obviously. The first thing to think that how bodybuilders learn to maintain BMI and musculoskeletal health.

Musculoskeletal health is also related with emotions which influence individual's suffering with muscle pain. People with musculoskeletal pain experiences pain with fibromyalgia (FM) which present elevated level of emotional distress (Crofford, 2015). Some positive and negative emotions considers some brain pathways to make connections with sensory regions which create pain (Bushnell, Ceko, Low. 2013).

A term psychogenic rheumatism was presented in a study which was diagnosed from various patients who had musculoskeletal pain that was also occurred fibrositis (a soft tissue rheumatic disorder) which can cause by environmental changes and can put extreme effects. Conversely, psychogenic rheumatism shows some environmental fluctuations which can lead emotions including mood, anxiety and diversions. These symptoms cause burning sensations, tightness, numbness, weakness and tingling sensations all the day. All these conditions impairs the ability of physical functioning. Psychotherapy is a primary treatment for this condition and physiotherapy is secondary treatment. (Barsky & Borus, 1999)

Methodology

The purpose of study was to check particular elements and to become aware of the bodily functions associated with hamstring muscle tightness by studying the bodily fitness and BMI of the bodybuilders including their emotional distress. To discover what fitness elements make contributions and impact on fitness of bodybuilders at gyms of Islamabad. A cross-sectional descriptive study was done and collected data from 377 body builders in Islamabad. The questionnaire included the demographic variables, private information and occupational traits. 126 people had been more than 25 years old, 50 people have been above 35 years. 30 people had been observed with extra weight problems. 82% people have been counted with regular BMI. About 150 participants were unmarried. Purposive sampling method was used. The demographic section of the questionnaire included Information approximately about their name, age, gender, occupation, weight, height, BMI, district, ethnicity, musculoskeletal disorder, any neurological situation and records of trauma. Three scales lower extremity functional scale (LEFS) and Musculoskeletal Health Questionnaire (MSHQ), Assessing emotional scale (AES) were used and data was analyzed with SPSS.

Results

Table: 1 Basic details of basic demographics of samples

Sr. No	Variables	Mean± SD
1	Name	1.00±.000
2	Age (years)	1.59±.667
3	Gender	1.00±.000
4	Occupation	1.99±.526
5	Weight (kg/m ²)	1.90±.600
6	Height (in feet's)	2.18±.529
7	BMI	1.76±.427
8	District	1.36±.658
9	Ethnicity	1.71±1.25
10	Muscular skeletal disorder	1.97±.169
11	Neurological condition	2.00±.000

12	History of trauma	2.00±.000
----	-------------------	-----------

Table: 2 Reliability of the lower extremity functional scale (LEFS) and Musculoskeletal Health Questionnaire and Assessing Emotional Scale (AES).

Scales	No. of items	Cronbach’s reliability co-efficient
LEFS	20	.99
MSHQ	14	.90
AES	31	.92

Table: 3 Mean, Standard Deviation, Reliability Coefficient of Scales (N=377).

Scales	N	M	SD	No. of items	Alpha coefficient
LEFS	377	64.76	9.18	20	.99
MSHQ	377	52.60	2.05	14	.90
AES	377	84.26	5.05	31	.92

Table 4: Prevalence of Hamstring Muscle Tightness according to Clinical Exposure.


Exposure	Normal hamstring	Tight hamstring	Toxic Emotions
Clinically exposed	220	50	275
Clinically non-exposed	157	65	102
Total	377	115	377

Figure 1: Bar Graph shows the percentage of bodybuilders classified among different emotional responses categories.




Bar graphs indicates emotional responses for maximum of as underscore 1 (24.7%), score 2 (37.3%), score 3 (12.5%), score 4 (5.5) and score 5 (20%) respectively. This bar graph indicates better percentage on y-axis approximately (24.7%) and decrease as (5.5%). This percentage of responses shows the responses of individuals according to the scoring key.

Figure 2: Bar graph shows the percentage of total score of the lower extremity Functional scale (LEFS).



The overall rating suggests the proportion of overall rating of bodybuilders that indicates the ratings 53 is the least rating and the best rating is seventy two this is closest to the reduce off rating 80. The bar graph display the 10% participants were given (53) rating, 20% (62), 30%(65), 40%(66), 50%(67), 60%(68), 70%(69), 80%(71) and 90% participants were given maximum rating of (72). The more the rating the decrease extremity. The decrease the rating the better extremity.

Figure 3: Bar graph shows the percentage of musculoskeletal health questionnaire



The overall rating suggests the share of general rating of bodybuilders that indicates the rating 50 is the least rating and the very best rating of this scale is 55. The decrease of rating is 56 of this scale that suggests the people were given better ranked ratings. Bar graph indicates 10% bodybuilders were

given (50) rating, 20 %(51), 40 %(52), 60 %(fifty three), 80 %(54) and 90% bodybuilders were given (55) rankings. The better the rating the more the musculoskeletal fitness.

Figure 4: Prevalence of Hamstring tightness (N=377)




Figure 4 shows the percentage of prevalence of hamstring tightness among bodybuilders on pie chart. This pie chart describes that the 82% of the bodybuilders with prevalence of hamstring tightness and the 18% with higher hamstring tightness.

Figure 5: percentage of toxic emotions and stable emotions.




Figure 5 suggests the share of incidence of hamstring tightness amongst bodybuilders on pie chart. This pie chart describes that the 82% of the bodybuilders with occurrence of hamstring tightness and the 18% with better hamstring tightness.

Figure 6: Relationship between hamstring tightness and higher emotional stability.




Figure 6 indicates the connection among the hamstring tightness with better emotions that describes the better emotional growth with the possibilities of occurrence of hamstring. This determine suggests approximately 82% of emotional values and 12% of hamstring tightness. As the emotional stability increases the hamstring tightness will decrease.

Figure 7: Relationship between hamstring tightness and lower emotional stability.




Figure 7 indicates the connection among the hamstring tightness with lower emotions that describes the possibilities of occurrence of hamstring. This determine suggests approximately 82% of hamstring values and 12% of lower emotional values. As the emotional stability decreases the hamstring tightness will increase.

Figure 8: Relationship between lower extremity function and musculoskeletal health and lower toxic emotions in prevalence of hamstring tightness.




Figure 8 suggests the connection among decrease extremity, toxic emotions and musculoskeletal fitness of hamstring tightness. This describes that better the ratings on decrease emotional 5 and extremity together with 72 and 55 on musculoskeletal fitness growth the superiority of hamstring tightness approximately 82%.

Discussion

The purpose to examine was to check the prevalence of hamstring tightness in bodybuilders of Islamabad who are having toxic emotions. It found out that the range of bodybuilders has excellent extremity and musculoskeletal fitness who have lower emotional control. Muscle tightness can be connected to emotions. Both can make contributions to more than one musculoskeletal conditions.

Muscle tightness may be linked to postural disturbances. Both can contribute to multiple musculoskeletal conditions (Spine, 2015). According to the Jose et al (2014), clinical observations suggested that hamstring tightness influences lumbar pelvic rhythm. Movement restrictions or postural asymmetry likely lead to compensatory movement patterns of the lumbar spine and subsequently to increased stress on the spinal soft tissues and increased risk of lower back pain (Esola, 1976).

The results of the study were emerged as relatively high prevalence of hamstring tightness among bodybuilders in Islamabad. The association between hamstring tightness, lower extremity functions and musculoskeletal health and emotions was significantly positively correlated at 0.05. It is determined that the prevalence of hamstring tightness among bodybuilders is very high (Bhagyashree, 2018).

An increase in hamstring flexibility was observed in all participants. The different studies were held among the different population about prevalence of hamstring tightness that showed the same results. A study conducted in

between the students of college colleges who showed 82% prevalence of hamstring tightness and the workers in offices showed prevalence about 85.7% (Nikolaos, 2012).

Due to lack of adequate physical activity, muscle weakness and some degenerative factors like osteoarthritis, senile osteoporosis and degenerative disk disease low back pain occurs in elderly people (Waqas, 2016). By feeling tense and stressed the individual may get achy areas, sore or downright painful areas. It occurs in shoulders, neck, hips and lower back. Stress can cause physical discomfort as the continuous weight of muscle contract.

Musculoskeletal problems in adults can sometimes be caused by differences in the shape of bones and joints that develop early in life. Muscular hypertension is caused by mental stress effort and bad attitude. Lack of blood flow on these muscle sites in the body can be disturbed.

The various researches were supported to the idea that the young people have higher tendencies to have greater level of prevalence of hamstring tightness and the men have lower tendency to have hamstring muscle flexibility (Wright et al, 1999). Physiotherapists should also include hamstring stretching exercises into the treatment programs, it may reduce the possible contribution of hamstring tightness to these disorders, especially low back pain syndrome (Fatima et al, 2017). And psychologist should also include some cognitive and behavior therapies for emotional prevalence. Deep breathing is best balance activity.

Clinical Implications

Stretching as soon as every week suggests exact end result with hamstring flexibility. Therefore habitual stretching of the hamstring muscular tissues want to be trained to all age groups, in particular earlier than age 30 while the tightness appears to grow greatly.

References

- Akinpelu, A.O., U. Bakare, and B.A. Adegoke, Influence of age on hamstring tightness in apparently healthy Nigerians. *Journal of Nigeria Society of Physiotherapy*, 2009. 15(2): p. 35-41
- Bhavana S, Mhatre, BS, Singh YL, Tembhekar JY, Mehta A, Which is the better method to improve “perceived hamstrings tightness” – Exercises targeting neural tissue mobility or exercises targeting hamstrings muscle extensibility? *Int J Osteopathic Med*, 2013
- Barsky AJ, Borus JF. Functional somatic syndromes. *Annals of internal medicine*. 1999;130:910. Doi: 7326/0003-4819-130-11-199906010-00016
- Bhagyashree KK & Deepak BA. “prevalence and severity of hamstring tightness among college students: A cross-sectional study.” 2018.
- Crofford. LJ. Psychological Aspects of chronic musculoskeletal pain, *Best Practice Research clinic Rheumatol*, 2015 May 21;29(1):147-155. Doi: 10.1016/j.berh.2015.04.027

- Dionne CE, Dunn KM, Croft PR, et al. A consensus approach toward the standardization of back pain definitions for the use in prevalence studies. *Spine (Phila Pa 1976)* 2008;33:95-103
- Erkula G, Demirkan F, Alper Kılıç B, Kiter E, “Hamstring shortening in healthy adults. *Journal of back and musculoskeletal rehabilitation,*” 2002.
- Esola MA, McClure PW, Fitzgerald GK, Siegler S. Analysis of lumbar spine and hip motion during forward bending in subjects with and without a history of low back pain. *Spine (Phila pa 1976)* 1996;21:71-78
- Fatima G, Qamar MM, Hassan JU, Basharat A, “Extended sitting can cause hamstring tightness. *Saudi Journal of Sports Medicine,*” 2017.
- Gleim GW, McHugh MP. Flexibility and its effects on sports injury and performance. *Sports Med.* 1997;24(5):289-299
- Koli BK, Anap DB, “. Prevalence and Severity of Hamstring Tightness among College Student: A Cross Sectional Study. *International Journal of Clinical and Biomedical Research.*” 2018.
- Mistry GS, Vyas NJ and Sheth MS. Correlation of hamstrings flexibility with age and gender in subjects having chronic low back pain. *International Journal of Therapies and Rehabilitation Research,* 2014. 3(4): p. 1.
- Mayorga-Vega D., et al., A physical education-based stretching program performed once a week also improves hamstring extensibility in schoolchildren: a cluster-randomized controlled trial. 2015.
- Nikolaos PT, Age-related differences of hamstring flexibility in male soccer players. *Baltic journal of health and physical activity,* 2012. 4(2): p. 110-115
- Raether PM, Lutter LD, “Recurrent compartment syndrome in the posterior thigh: report of a case. *The American journal of sports medicine,*” 1982.
- Shah AJ, Iftakhar MT, “Cause and Management of Hamstring Injuries in College Sprinters-A Qualitative Approach. *J Sports Med Doping Stud,*” 2017
- Spine J. (2015), influence of hamstring tightness in pelvic, lumbar and trunk range of motion in the low back pain and asymptomatic volunteers during forward bending.
- Sandeep S, et al, “effect of mobilization and PNF stretching on hamstring Flexibility in working women”, 2015.
- Weerasekara I, Kumari I, Weerathna N, Withanage C, Wanniarachchi C, “The prevalence of hamstring tightness among the male athletes of University of Peradeniya in 2010, Sri Lanka. *Int J Phys Med Rehab*” 2013
- Wright, G, A., Delong, T. H., & Gehlsen, G. “Electromyographic activity of the hamstring during performance of the leg curl, stiff-leg deadlift, and back squat movements. *The journal of strength & conditioning research,* 1999
- Waqas MS, Naqvi SMA, Hussain HS, et al. “Frequency of reduced hamstring flexibility in prolonged sitting (6-8 hours) among office workers. *Journal of Riphah College of Rehabilitation sciences*” 2016.